

Emil Mottola, T-2, Los Alamos National Laboratory

Scalar Gravitational Waves in the Effective Theory of Gravity

Abstract: As a low energy effective field theory, classical General Relativity receives an infrared relevant modification from the conformal trace anomaly of the energy-momentum tensor of massless, or nearly massless, quantum fields. The local form of the effective action associated with the trace anomaly is expressed in terms of a dynamical scalar conformal field that couples to the conformal factor of the spacetime metric, allowing it to propagate over macroscopic distances. Linearized around flat spacetime, this semi-classical EFT admits scalar gravitational wave solutions in addition to the transversely polarized tensor waves of the classical Einstein theory. The strongest astrophysical sources for scalar gravitational wave burst signals detectable by LIGO and VIRGO are the excited gluonic condensates in the interiors of neutron stars as they are formed or merge with other compact objects. Primordial scalar gravitational waves should also have been copiously generated by the hot QCD plasma in the early universe, whose detection likely requires space-based GW antennae.

Biography: B. A. Astrophysics 1974 Columbia College, NY

Ph. D. Physics 1979 Columbia Univ. NY

Post-doctoral Member, Institute for Advanced Study, Princeton, NJ 1979-1982

Post-doctoral Research: Institute for Theoretical Physics (now KITP), UCSB, Santa Barbara, CA 1982-1986

Staff Member, LANL Elementary Particles & Field Theory, T-8, 1986-2008

Deputy Group Leader, LANL T-8, 1995-2000

Scientific Associate, CERN Theory Div. 2009-2010

Staff Member, LANL, Theoretical Division, Nuclear, Particle, Astrophysics & Cosmology, T-2, 2008-present

Approximately 100 Publications, 5000+ Citations